

# IV Q6 BUSULFAN PHARMACOKINETICS REQUISITION

## PATIENT INFORMATION

Patient Name: \_\_\_\_\_  
 Medical Record #: \_\_\_\_\_  
 Actual Weight (kg): \_\_\_\_\_  
 Dosing Weight (kg): \_\_\_\_\_  
 Height (cm): \_\_\_\_\_

Full Institution Name: \_\_\_\_\_  
 Date of Birth: \_\_\_\_\_  
 Genetic Sex (check one): Male / Female  
 Diagnosis: \_\_\_\_\_  
 ICD-10 Code: \_\_\_\_\_

## DOSE INFORMATION

Date of Dose: \_\_\_\_\_  
 Dose Given (mg): \_\_\_\_\_  
 Busulfan Manufacturer/Lot Number  
 (if generic busulfan): \_\_\_\_\_  
 Dose Number \_\_\_\_\_ Total # of  
 (write "test" for a test dose): \_\_\_\_\_ Regimen doses: \_\_\_\_\_  
 Desired Target Range: \_\_\_\_\_  
 Target Units (circle one):  $\mu\text{Mol}\cdot\text{min}$  /  $\text{mg}\cdot\text{h/L}$  /  $\text{ng/mL}$   
(AUC) (AUC) (Css)  
 For a test dose, indicate the regimen  
 dosing interval (circle one): Q6 / Q8 / Q12 / Q24

## CONTACT INFORMATION

Signature of MD or designee: \_\_\_\_\_  
 Attending MD (print name): \_\_\_\_\_  
 Results are usually available between 13:00 and 16:00 Pacific Time the day following  
 sample collection and shipment. Verbal report recipient must be an MD or a PharmD.  
 Verbal report recipient: \_\_\_\_\_  
 Verbal report recipient  
 contact number: \_\_\_\_\_  
 Email address(es)/Fax number(s): \_\_\_\_\_

**ALL INFORMATION MUST BE FILLED OUT PRIOR TO SHIPPING.**

**BUSULFAN RESULTS CANNOT BE CALCULATED OR REPORTED WITHOUT COMPLETE INFORMATION.**

### IV Q6 Busulfan Dose 1 or Test Dose

Typical IV Q6 or Test Dose infusions are 120 minutes, including flush

Infusion start time: \_\_\_\_\_  
 Infusion stop time: \_\_\_\_\_

	ACTUAL Sample Collection Clock Times	Initials
End of Infusion		
End of Infusion + 15 Minutes		
End of Infusion + 30 Minutes		
Start of infusion + 4 Hours		
Start of infusion + 5 Hours		
Start of infusion + 6 Hours		

### IV Q6 Busulfan Follow-up Doses

Typical IV Q6 infusions are 120 minutes, including flush

Infusion start time: \_\_\_\_\_  
 Infusion stop time: \_\_\_\_\_

	ACTUAL Sample Collection Clock Times	Initials
<b>Pre Infusion:</b>		
End of Infusion		
End of Infusion + 15 Minutes		
Start of infusion + 4 Hours		
Start of infusion + 5 Hours		
Start of infusion + 6 Hours		

Please draw a minimum of 2 mL blood in a green top tube (sodium heparin). Keep refrigerated or on ice at all times. Centrifuge at 4°C. Remove and freeze plasma into a plastic tube labeled with: Patient Name, Medical Record #, Date and Time of Draw. Please tape labels on. Send plasma with 5 kg of dry ice **FIRST OVERNIGHT** to the address below. Accurate blood draw and infusion start/stop times are critical to busulfan PK analysis.

#### DRUG INTERACTIONS: Please indicate which (if any) of the following drugs the patient has taken within the past 30 days:

Deferasirox, Metronidazole, Itraconazole, Isavuconazole, Voriconazole, Posaconazole, Azithromycin, TKIs, Acetaminophen, Ivosidenib, Enasidenib

Drug(s): \_\_\_\_\_  
 Date of last dose: \_\_\_\_\_

#### Please indicate any other drug/treatment the patient has taken or will take as part of their current conditioning regimen:

Cyclophosphamide	Thiotepa	Etoposide
Fludarabine	ATG	TBI
Melphalan	Other: _____	

**Please fax or scan and email a completed copy of this requisition form and shipment tracking number to PKLab@seattlecca.org prior to shipping samples, and include a hard copy with the samples. Ship samples frozen with a minimum of 5kg dry ice.**

**Phone number: (206) 606-7389      Fax number: (206) 606-7397      Pager: (206) 994-5942**  
**Email: PKLab@seattlecca.org**

**SHIP TO:** Pharmacokinetics  
 Laboratory  
 Seattle Cancer Care Alliance  
 188 E. Blaine St. Suite 250  
 Seattle, WA 98102