SCCA Clinic SARS-CoV-2 (COVID-19) and Respiratory Viral Testing Guidelines for Patients

These guidelines are intended to provide guidance on testing for SARS-CoV-2 and other respiratory viruses among immunocompromised patients. As you are all aware, this is a rapidly evolving situation and testing guidance will be updated accordingly depending on community transmission dynamics and test capacity. Do not ignore other signs and symptoms during this COVID-19 period, as patients with fever and signs of infection should be treated by current guidelines (e.g. neutropenic fever, sepsis) by inpatient services.

Clinical Criteria	SARS-CoV-2 PCR Test?	Extended Respiratory Viral PCR Test?	Infection Control Precautions
Asymptomatic (absence of any of the below symptoms)	BMT/ IMTX (weekly screening); pre-aerosol generating procedure; pre-radiation	NO	Standard Precautions
No respiratory symptoms <u>BUT</u> Fever or other symptoms (anosmia, dysgeusia, new-onset diarrhea)	YES	NO	Droplet + Contact Precautions (+/- Enteric Precautions)
Respiratory symptoms*	YES – • All SCCA Patients	 YES – Extended Respiratory Viral PCR Test Only BMT/ IMTX, hematologic malignancy 	Droplet + Contact Precautions
Chest CT with new ground glass opacities WITH OR WITHOUT symptoms	• All SCCA Patients	 YES – Extended Respiratory Viral PCR Test Only BMT/ IMTX, hematologic malignancy 	Droplet + Contact Precautions

Refer to Infection Prevention Guidelines for removing patients from precautions.

^{*}Respiratory symptoms: cough, shortness of breath, wheezing, sore throat, congestion, runny nose, myalgias